Oral Communication 7

The effect of synthetic cannabinoid ADB-FUBI-NACA on primary neuronal cultures ß-galactosidase activity: preliminary findings

<u>Rita Roque-Bravo</u>^{1,2,*}, Helena Carmo^{1,2}, João Pedro Silva^{1,2}, Félix Carvalho^{1,2} and Diana Diasda-Silva^{1,2,3,4,5,*}

¹ UCIBIO, Laboratory of Toxicology, Department of Biological Sciences, Faculty of Pharmacy, University of Porto, Rua Jorge Viterbo Ferreira 228, 4050-303 Porto, Portugal

² Associated Laboratory i4HB – Institute for Health and Bioeconomy, Faculty of Pharmacy, University of Porto, Rua Jorge Viterbo Ferreira 228, 4050-303 Porto, Portugal

³ LAQV/REQUIMTE, ESS, Polytechnic of Porto, Rua Dr. António Bernardino de Almeida, 400 4200 - 072, Porto, Portugal ⁴ Associate Laboratory i4HB – Institute for Health and Bioeconomy, University Institute of Health Sciences – CESPU, 4585-116 Gandra, Portugal

⁵ UCIBIO – Applied Molecular Biosciences Unit, Forensics and Biomedical Sciences Research Laboratory, University Institute of Health Sciences (1H-TOXRUN, IUCS-CESPU), 4585-116 Gandra, Portugal

* Correspondence: rita.rdsb@gmail.com (RRB); dds@ess.ipp.pt (DDS)

Abstract

Background: ADB-FUBINACA (ADB-FUB) is a synthetic cannabinoid (SC) that has gained popularity among users as a new psychoactive substance. This stems from SC's pharmacological similarity to the active principle of cannabis, Δ^9 -tetrahydrocannabinol (THC). However, unlike THC, SCs demonstrate full agonism of cannabinoid receptors 1 and 2 [1]. Recent scientific developments have shown that cannabis use may aggravate ageing-related parameters [2,3]. Moreover, a study using human fibroblasts revealed that 1 µM THC (2h-long exposure, for 15 days) can increase β-galactosidase activity [3], which serves as a first-line marker for cellular senescence. Objective: This study was designed to investigate whether these biologically-relevant concentrations could accelerate neuronal ageing. Methods: PHC were isolated from Wistar rat day 18-19 embryos and cultured for up to 21 days-in-vitro (DIV). Exposure to 1 pM, 1 nM and 1 µM ADB-FUB (concentrations previously shown to be non-cytotoxic to PHC) started either on DIV3 or DIV7 and was maintained up to 21 DIV. At that final timepoint, ß-galactosidase activity was evaluated. DMSO at 0.02% was employed as solvent control. Results: Under these experimental conditions, PHC exposed to 1 nM and 1 µM ADB-FUB in the DIV3-21 protocol had lower ßgalactosidase activity when compared to control conditions (p<0.05, 1 nM; p<0.001, 1 μ M). No statistically significant results were registered for PHC under the DIV7-21 exposure protocol. Conclusions: These findings are, to the best of our knowledge, the first evidence of a potential "anti-ageing" effect of ADB-FUB. Evaluation of other senescence-related endpoints will follow. Moreover, experiments using another in vitro neuronal model (human neuroblastoma cell line SH-SY5Y) are underway to compare the effects of the same drug in different models and further substantiate conclusions on ADB-FUB's effect.

Keywords: synthetic cannabinoids; in vitro neurotoxicity; ageing

Acknowledgments

This work was supported by FCT—Fundação para a Ciência e a Tecnologia (projects UIDP/04378/2020 and UIDB/04378/2020 of the Applied Molecular Biosciences—UCIBIO), the project LA/P/0140/2020 of the Associate Laboratory Institute for Health and Bioeconomy—i4HB, the PhD grant 2020.04493.BD (RRB) and research contract (under Scientific Employment Stimulus) 2021.01789.CEECIND/CP1662/CT0014 (JPS).

References

- 1. Kemp et al. Top 10 facts you need to know about synthetic cannabinoids: not so nice Spice. Am. J. Med. 2016; 129: 240-44.el.
- Burggren, A.C.; Siddarth, P. Subregional Hippocampal Thickness Abnormalities in Older Adults with a History of Heavy Cannabis Use. *Cannabis Cannabinoid Res.* 2018; 3(1): 242-51.
- Allen, J.P.; Danoff, J.S. Lifetime marijuana use and epigenetic age acceleration: A 17-year prospective examination. *Drug Alcohol Depend*. 2022; 233, 109363.
- 4. Gerasymchuk, M.; Robinson, G.I. Phytocannabinoids Stimulate Rejuvenation and Prevent Cellular Senescence in Human Dermal Fibroblasts. *Cells*. 2022; 11: 3939.



In *Scientific Letters*, works are published under a CC-BY license (Creative Commons Attribution 4.0 International License at https://creativecommons.org/licenses/by/4.0/), the most open license available. The users can share (copy and redistribute the material in any medium or format) and adapt (remix, transform, and build upon the material for any purpose, even commercially), as long as they give appropriate credit, provide a link to the license, and indicate if changes were made (read the full text of the license terms and conditions of use at https://creativecommons.org/licenses/by/4.0/).